



Curvature profiles from stochastic inflation


COSMOLOGICAL PROBES OF THE EARLY UNIVERSE

1 JULY 2026, KING'S COLLEGE LONDON

EEMELI TOMBERG, UCLouvain


Primordial perturbations

Long scales: weak perturbations
CMB, large-scale structures



Primordial perturbations


Long scales: weak perturbations
CMB, large-scale structures



Short scales: ?
Primordial black holes?



Primordial black holes




Dark matter

Seeds of supermassive BHs

GW source


Primordial black holes




Zel'dovich and Novikov 1967

Hawking 1971 and Carr 1974


Date of paper




PBHs from inflation



PBHs from inflation



Longer perturbations $\leftrightarrow$ larger PBH mass



Which perturbations collapse?


Numerical relativity
simulations

Compaction function:

$$\mathcal{C}(r, t) = 2 \frac{\delta M(r, t)}{R(r, t)}$$

Collapse:

$$\mathcal{C}_{\max} > \mathcal{C}_{\text{th}} = \delta_c$$



Compaction versus curvature

Super-Hubble:

$$\mathcal{C}(r) = \frac{2}{3} (1 - [1 + r\zeta'(r)]^2)$$

Note assumption of spherical symmetry

Inflationary perturbations

Background

Equations of motion:

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0, \quad 3H^2 = \frac{1}{2}\dot{\phi}^2 + V(\phi)$$

Slow-roll parameters:

$$\epsilon_1 \equiv -\partial_N \ln H, \quad \epsilon_2 \equiv \partial_N \ln \epsilon_1$$



Inflation?



Slow-roll?

Perturbations

Field perturbations:

$$\delta\ddot{\phi} + 3H\delta\dot{\phi} + \left[\frac{k^2}{a^2} + V''(\phi) - \frac{1}{a^3} \frac{d}{dt} \left(\frac{a^3}{H} \dot{\phi}^2 \right) \right] \delta\phi = 0$$


Curvature perturbation:

$$\zeta_k = \frac{\delta\phi_k}{\sqrt{2\epsilon_1}} \sim H$$

$$\mathcal{P}_\zeta(k) = \frac{k^3}{2\pi^2} |\zeta_k|^2$$

Beyond linear perturbations

Separate universe approximation:



ΔN approximation: $\Delta N = N - \langle N \rangle = \zeta$

Divide the field in two

$$\phi_{\text{tot}} = \phi + \delta\phi$$


$$\int_{k < k_\sigma} \frac{d^3k}{(2\pi)^{2/3}} \phi_k(t) e^{-i\vec{k} \cdot \vec{x}}$$


Separate universe


$$\int_{k > k_\sigma} \frac{d^3k}{(2\pi)^{2/3}} \phi_k(t) e^{-i\vec{k} \cdot \vec{x}}$$


Linear perturbations

Coarse-graining scale: $k = k_\sigma \equiv \sigma a H$


Drifting modes: stochasticity



Drifting modes: stochasticity



Drifting modes: stochasticity



Stochastic inflation

$$\partial_N \phi = \pi + \xi_\phi, \quad \partial_N \pi = -\left(3 - \frac{1}{2}\pi^2\right) \left(\pi + \frac{V'(\phi)}{V(\phi)}\right) + \xi_\pi$$

e-folds as time variable

$$\langle \xi_X(N) \xi_Y(\tilde{N}) \rangle = \mathcal{P}_{XY}(N, k_\sigma) \delta(N - \tilde{N})$$



de Sitter result / numerical computation


Stochastic inflation

Slow roll:

$$\partial_N \phi = -\frac{V'(\phi)}{V(\phi)} + \xi_\phi$$

$$3H^2(\phi) = V(\phi)$$

$$\langle \xi_\phi(N) \xi_\phi(\tilde{N}) \rangle = \frac{H^2(\phi)}{4\pi^2} \delta(N - \tilde{N})$$



Radial profiles

Raatikainen:2023bzk

$$\begin{aligned}\Delta N_{<k} &= N_{<k} - \langle N_{<k} \rangle \\ &= \zeta_{<k}(\mathbf{x}) \equiv \int \frac{d^3p}{(2\pi)^{3/2}} \zeta_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} \theta(k - p)\end{aligned}$$

Radial profiles

Raatikainen:2023bzk

$$\begin{aligned}\Delta N_{<k} &= N_{<k} - \langle N_{<k} \rangle \\ &= \zeta_{<k}(\mathbf{x}) \equiv \int \frac{d^3p}{(2\pi)^{3/2}} \zeta_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} \theta(k - p)\end{aligned}$$

Vary coarse-graining scale for same patch

Radial profiles

Raatikainen:2023bzk



$$\begin{aligned}\Delta N_{<k} &= N_{<k} - \langle N_{<k} \rangle \\ &= \zeta_{<k}(\mathbf{x}) \equiv \int \frac{d^3p}{(2\pi)^{3/2}} \zeta_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} \theta(k - p)\end{aligned}$$

Vary coarse-graining scale for same patch


$$\text{Obtain: } \zeta_k = \sqrt{\frac{\pi}{2}} \frac{1}{k^3} \frac{d\zeta_{<k}}{d \ln k} \rightarrow \zeta(r) = \int_0^\infty \frac{dk}{k} \frac{d\zeta_{<k}}{d \ln k} \frac{\sin(kr)}{kr}$$

Primordial black hole models


Inflection point models




Inflection point models



Exponential tails



Simplification: constant roll



ϵ_2 approx. constant


$$\partial_N \phi = \frac{\epsilon_2}{2} \phi + \xi_\phi$$



$$\langle \xi_\phi(N) \xi_\phi(\tilde{N}) \rangle = \mathcal{P}_\phi(N, k_\sigma) \delta(N - \tilde{N})$$

Tomberg:2023kli



Simplification: constant roll

Tomberg:2023kli







Raatikainen:2025gpd




Raatikainen:2025gpd




EXTREMELY SPIKY


Raatikainen:2025gpd




$P(C)$ $P(C_T)$ $P(C_{\text{smooth fit}})$
 $P(\bar{C})$ $P(\bar{C}_T)$ $P(\bar{C}_{\text{smooth fit}})$





$P(C)$ $P(C_{\text{mean}})$ $P(C_{\text{Gauss mean}})$
 $P(\bar{C})$ $P(\bar{C}_{\text{mean}})$ $P(\bar{C}_{\text{Gauss mean}})$






Raatikainen:2025gpd





Raatikainen:2025gpd

- No critical scaling
- - - Reverse lognormal fit
- Critical scaling
- Transfer function
- Transfer function with critical scaling



Open questions

Behaviour at large radius?

Correct collapse criterion
for spiky profiles?

Redo collapse simulations?

SUMMARY

Primordial black hole predictions require non-linear techniques: stochastic inflation

To compute the compaction function,
need curvature perturbation profile

Stochastic compaction function profiles
are very spiky

$$\ddot{\phi} - \frac{1}{a^2} \nabla^2 \phi + 3H\dot{\phi} + V'(\phi) = 0$$

Linear perturbations:
small- $\delta\phi$ expansion

$$\ddot{\phi} - \frac{1}{a^2} \nabla^2 \phi + 3H\dot{\phi} + \underline{V'(\phi)} = 0$$

$\delta\ddot{\phi}$ $\delta\phi$ $\delta\dot{\phi}$ $V''(\phi)\delta\phi$ + metric perts


Separate universe:
small- $\frac{\nabla^2}{a^2 H^2}$ expansion

$$\ddot{\phi} - \frac{1}{a^2} \nabla^2 \phi + 3H\dot{\phi} + V'(\phi) = 0$$

Stochastic inflation

“Stochastic inflation” in
INSPIRE-HEP

- 164 papers
- 98 of these since 2015




Stochastic trees

Animali:2025pyf

Start from 1 Hubble patch

Once volume doubled: split in two

Spatial distribution?
Clustering?



Lattice simulations

Mizuguchi:2024kbl

Stochastic grid

Launay:2024qsm

Combining stochastics and numerical relativity

PREPARED FOR SUBMISSION TO JCAP

RUP-24-10

STOLAS: STOchastic LAttice Simulation of cosmic inflation



Yurino Mizuguchi,^a Tomoaki Murata,^b and Yuichiro Tada^{c,a}

^aDepartment of Physics, Nagoya University,
Furo-cho Chikusa-ku, Nagoya 464-8602, Japan

^bDepartment of Physics, Rikkyo University,
Toshima, Tokyo 171-8501, Japan

^cInstitute for Advanced Research, Nagoya University

Stochastic Inflation in General Relativity

Yoann L. Launay,^{1,*} Gerasimos I. Rigopoulos,^{2,†} and E. Paul S. Shellard^{1,‡}

¹*Centre for Theoretical Cosmology, Department of Applied Mathematics and Theoretical Physics,
University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, United Kingdom*

²*School of Mathematics, Statistics and Physics, Newcastle University,
Newcastle upon Tyne, NE1 7RU, United Kingdom*

We provide a formulation of Stochastic Inflation in full general relativity that goes beyond the slow-roll and separate universe approximations. We show how gauge invariant Langevin source terms can be obtained for the complete set of Einstein equations in their ADM formulation by providing a recipe for coarse-graining the spacetime in any small gauge. These stochastic source terms are defined in terms of the only dynamical scalar degree of freedom in single-field inflation and all depend simply on the first two time derivatives of the coarse-graining window function, on the gauge-invariant mode functions that satisfy the Mukhanov-Sasaki evolution equation, and on the slow-roll parameters. It is shown that this reasoning can also be applied to include gravitons as stochastic sources, thus enabling the study of all relevant degrees of freedom of general relativity for inflation. We validate the efficacy of these Langevin dynamics directly using an example in uniform field gauge, obtaining the stochastic e-fold number in the long wavelength limit without the need for a first-passage-time analysis. As well as investigating the most commonly used gauges in cosmological perturbation theory, we also derive stochastic source terms for the coarse-grained BSSN formulation of Einstein's equations, which enables a well-posed implementation for 3+1 numerical relativity simulations.

I. INTRODUCTION

Inflation theory was postulated more than 40 years ago as an explanation for the apparently fine-tuned initial conditions of the Hot Big Bang [1–3]. The proposal gained traction as it also offers a natural mechanism for generating the initial density inhomogeneities [4–9] which in later stages of cosmic history led to the formation of cosmic structure via gravitational instability. These den-


efforts [13–18], there are regimes where predictions can still be made by using techniques from Quantum Field Theory on Curved Spacetime (QFTCS) [15] or by constructing Effective Field Theories (EFTs), which have made continuous advancements in cosmology, inspired by the latter's success in flat space [19]. Abandoning pretenses of completeness, an EFT establishes a region of validity, normally bounded by ultraviolet (UV) and/or infrared (IR) cutoffs, and the narrative of theoretical physics is implicitly about pushing these cutoffs to their

[astro-ph.CO] 20 Dec 2024

30v2 [gr-qc] 7 May 2024

Multi-field inflation

Murata:2025onc





Hybrid inflation model

Upper bound to curvature
perturbation

Gradient corrections

Briaud:2025ayt

Go beyond leading order in gradient expansion




Colored noise
Exponential tails: a pullback effect

Eternal inflation

Tomberg:2025fku

$$\text{eternal inflation} \iff \lim_{N \rightarrow \infty} \langle V \rangle_N \equiv \lim_{N \rightarrow \infty} \int_{\phi_{b1}}^{\phi_{b2}} e^{3N} P(\phi, N) d\phi > 0$$



$$P(\phi, N) \sim e^{-\lambda N} \quad \lambda \leq 3$$

$$\text{Maximum:} \\ \lambda \approx |\eta_H| \sim 0.1$$

$$\text{Minimum:} \\ \lambda \approx 0$$

Eternal inflation

